



# Management of In-Flight Medical Emergencies

This supplementary material has been provided by the authors to give readers additional information about their work.

Martin-Gill C, Doyle TJ, Yealy DM. In-flight medical emergencies: a review. *JAMA*. doi:10.1001/jama.2018.19842





## SYNCOPE / NEAR-SYNCOPE

— 30% of all in-flight emergencies —

### Initial assessment-suspect

- Vasovagal:** Pale, diaphoretic, improves with simple measures in 15-30 min.
- Cardiac cause (eg, myocardial infarction):** Chest pain, dyspnea, arm or jaw pain, persistent bradycardia.
- Pulmonary:** Dyspnea, pleuritic chest pain.
- Stroke:** Slurred speech, facial droop, or arm weakness.
- Hypoglycemia:** Diaphoretic, cool skin; assess with glucometer if available.

### Management and expected course

- If unconscious** ▶ Lie flat, elevate legs, apply oxygen. If no pulse or signs of life, follow cardiac arrest card.
- If transient syncope** ▶ Supine position, elevate legs. Oral fluids with head raised if nausea absent. If improves in 15-30 min, slowly sit up and return to seat if tolerated.
- If hypoglycemia** ▶ Oral glucose or 25 g of dextrose 50% intravenously.
- If other conditions suspected** ▶ Refer to relevant card.
- If no improvement or not progressing as expected** ▶ Contact ground-based medical support for additional recommendations.



## GASTROINTESTINAL ILLNESS

— 15% of all in-flight emergencies —

### Initial assessment

- Identify extent and timing of symptoms, including nausea, vomiting, diarrhea, bleeding, and specifics of any abdominal pain (location, quality, and severity).

### Management and expected course

- If nausea/emesis** ▶ Use an oral antiemetic if available; if not tolerated, consider a parenteral antiemetic.
  - Provide oral hydration if tolerated.
  - Use sugar-containing liquids if symptoms of hypoglycemia.
  - If oral intake not tolerated, consider intravenous fluids.
- If dyspepsia** ▶ Use an antacid if available in the emergency medical kit.
- If diarrhea** ▶ Use an antidiarrheal if available in the emergency medical kit.
  - If patient has fever and persistent diarrhea (>14 d), contact ground-based medical support, as local public health authorities may need to be contacted at the destination.
- If severe abdominal pain, tenderness on examination, rigid abdomen, or blood in bodily fluid** ▶ Contact ground-based medical support for additional recommendations.



## RESPIRATORY DISTRESS

10% of all in-flight emergencies

### Initial assessment

- Identify history of respiratory disease, scuba diving, extremity swelling, or infectious symptoms.
- If available, check pulse oximetry.

### Management and expected course

- If ongoing dyspnea or known oxygen saturation is <95%** ▶ Administer oxygen.
  - If passenger's portable oxygen concentrator fails or is not used for a patient with preexisting lung disease, consider trial of oxygen therapy.
  - If passenger uses  $\geq 4$  L/min on the ground, the onboard oxygen supply may not be enough to reverse hypoxia.
  - Monitor flow rate of oxygen administered; canister consumption is variable and aircraft may not have sufficient oxygen for continuous use for the duration of the flight.
- If bronchospasm** ▶ Administer albuterol, 2.5 mg inhaled.
- If allergic reaction** ▶ Refer to allergic reaction card.
- If passenger does not improve** ▶ Contact ground-based medical support for additional recommendations.



## CARDIOVASCULAR SYMPTOMS

7% of all in-flight emergencies

### Initial assessment

- Identify if any prior myocardial infarction or other cardiovascular history.
- In some settings, a 12-lead electrocardiogram may be obtained and transmitted for ground review (and/or volunteer review if qualified to read).
- Suspected acute coronary syndrome:** Chest pain, dyspnea, arm or jaw pain.
- Suspected arrhythmia:** Persistent bradycardia, tachycardia, or irregular heartbeat.
- Suspected dyspepsia:** Isolated epigastric burning with no associated symptoms. This is a consideration of exclusion, supported by history of similar symptoms.

### Management and expected course

- If suspected acute coronary syndrome** ▶ Aspirin, 325 mg orally; nitroglycerin, 0.4 mg sublingually every 5-10 min (if systolic blood pressure is  $\geq 100$  mm Hg).
- If any dyspnea or respiratory distress** ▶ Give oxygen, unless saturations are known to be near or at normal levels.
- If dyspepsia suspected** ▶ Antacids or other analgesics can be given after appropriate risk stratification. Alternative causes should first be considered.
- If persistent or additional symptoms** ▶ Contact ground-based medical support for additional recommendations.



## STROKE-LIKE SYMPTOMS

Up to 5% of all in-flight emergencies

### Initial assessment

- A focused history should include the time of symptom onset, specific motor and sensory components, and any other associated symptoms including headache or sensorium changes.
- Screening for stroke:** Speech disturbance, facial droop, or arm weakness.

### Management and expected course

- Administer oxygen, unless saturations are known to be near or at normal levels.
- If patient has ongoing neurological deficits suggestive of a stroke** ▶ Contact ground-based medical support.
  - Recommendation may include diversion, which may not be to the closest airport if stroke care is not present at that airport.
  - Ground-based team should have information on capabilities for medical care near most major airports.



## SEIZURE

Up to 5% of all in-flight emergencies

### Initial assessment

- Identify the symptoms the passenger exhibited during the event:**

Including onset, duration of movement activity, quality of movements (eg, tonic-clonic), and loss of bowel or bladder function.

### Management and expected course

- If unresponsive** ▶ Lay passenger on floor on side, monitor airway, and assess vital signs with ongoing neurological examination as above.
- If ongoing seizing** ▶ Administer parenteral benzodiazepines if available in the emergency medical kit (not usually available on US commercial airlines).
- If alert following a prolonged or recurrent seizure** ▶ Ground-based medical support may recommend an added dose of the patient's own antiepileptic medication (if history of seizures and available) or an oral benzodiazepine (if available in the emergency medical kit).
- If seizure resolves and patient regains normal mental status** ▶ Diversion is not commonly necessary.



## TRAUMA

5% of all in-flight emergencies

### Initial assessment

- Assess all injuries for any open wounds, tenderness, deformity, or active bleeding.
- Assess patients with injury to the head, neck, or back for any neurological symptoms.

### Management and expected course

- Injuries from falling luggage** ▶ Typically minor and may be assessed further at the destination.
- Active bleeding** ▶ Control bleeding with direct pressure using a gloved hand.
- Ongoing heavy extremity bleeding** ▶ Consider applying a tourniquet.
- Suspected long bone or joint injuries** ▶ Splinting material is not commonly found in the emergency medical kit, but may be improvised from available equipment (eg, a U-shaped half-rolled magazine secured with tape will make a good forearm or wrist splint).





## PSYCHIATRIC SYMPTOMS

Up to 3% of all in-flight emergencies

### Initial assessment

- Aim to create a rapport with the passenger to deescalate the situation.
- Elicit information and consider the passenger's use of mood-altering substances.
- Identify if patient takes specific psychiatric medications, dosing, last dose taken, and if available on aircraft.

### Management and expected course

- If verbal deescalation ineffective** ▶ Consider a benzodiazepine if available from an extended emergency medical kit.
  - Benzodiazepines are not commonly available in the emergency medical kit and are infrequently necessary even when available.
- If combative** ▶ Refer to flight crew for individual airline security protocols, which take precedence over attempts at medical management.
  - Airline security protocols vary by airline and may include restraining the passenger or diverting the aircraft for the safety of other passengers and crew.



— Up to 3% of all in-flight emergencies —

### Initial assessment

- Identify type, amount, and timing of substances used.
- Identify symptoms and mental status, along with vital signs.
- Suspected opioid ingestion:** Altered mentation, constricted pupils, respiratory depression.
- Suspected alcohol ingestion:** Altered mentation, slurred speech, behavior changes.
- Suspected stimulant ingestion:** Altered mentation, tachycardia, dilated pupils, agitation.

### Management and expected course

- If normal vital signs and no respiratory compromise** ▶ Observation only.
- If suspected opioid ingestion with respiratory depression** ▶ Naloxone, 0.4-0.8 mg intravenously or 2 mg intramuscularly/intranasally.
- If suspected alcohol overdose** ▶ Observe and provide antiemetic therapy.
- If suspected stimulant ingestion** ▶ Observe and hydrate (for tachycardia). Consider benzodiazepine if available from the emergency medical kit.
- If ongoing respiratory distress or combativeness** ▶ Contact ground-based medical support for additional recommendations. Refer to airline crew for individual airline security protocols.



## ALLERGIC REACTION

2% of all in-flight emergencies

### Initial assessment

- Identify any known or likely allergen exposure; duration and severity of symptoms; and any airway swelling, respiratory involvement, or signs of systemic reaction such as generalized hives.
- Suspected local allergic reaction:** Localized pruritic rash or isolated hives.
- Suspected anaphylaxis:** Airway swelling, respiratory distress, generalized hives, hypotension, nausea/vomiting.

### Management and expected course

- If local allergic reaction ▶**  
Diphenhydramine, 25-50 mg in adults or 1 mg/kg in children orally.
  - If unable to tolerate oral ingestion, diphenhydramine intravenously/intramuscularly at above dose.
  - Try a different histamine blocker if available in the emergency medical kit.
- If anaphylaxis ▶** Epinephrine, 1 mg/mL (0.3 mL in adults, or 0.15 mL in children intramuscularly), diphenhydramine, and steroids if available in the emergency medical kit. Epinephrine may be available as an autoinjector or in an ampoule to be drawn up via syringe.
- If there is no improvement ▶**  
Contact ground-based medical support for additional recommendations.



1% of all in-flight emergencies

## Initial assessment

- Identify onset and detailed description of symptoms, along with information about the pregnancy (eg, parity, gestational age, and any preceding complications).
- Vaginal bleeding:** Assess duration and severity (ie, equivalent of pads per h).
- Labor suspected:** Regular contraction, gush of vaginal fluid.

## Management and expected course

- If vaginal bleeding <1 pad per h ▶**  
Expectant management is common.
- If preterm labor in third trimester ▶**  
Place the passenger on left side and consider fluid intravenously if any concerns exist for blood loss or distress.
- Active labor, ongoing/severe vaginal bleeding, or increasing/severe abdominal pain ▶**  
Contact ground-based medical support for additional recommendations.



## CARDIAC ARREST

0.2% of all in-flight emergencies

### Initial assessment

- Check breathing and pulse; limit pulse checks to <10 seconds.

### Management and expected course

- If no pulse or signs of life ▶**
  - Start chest compression-only cardiopulmonary resuscitation, with addition of bag-valve-mask ventilation (30 compressions to 2 ventilations) when the emergency medical kit is available and someone skilled is present.
  - Obtain and apply automated external defibrillator as soon as possible and follow instructions for defibrillation.
  - If no shock is advised, or AFTER a shock is delivered, resume cardiopulmonary resuscitation if there is no pulse.
  - If no response to cardiopulmonary resuscitation and automated external defibrillator, initiate an intravenous line. Administer epinephrine (0.1 mg/mL) 1 mg intravenously, along with consideration of causal reversible conditions such as hypovolemia and tension pneumothorax.
- Instruct flight crew to notify the ground team and pilot if not already done. If no shock is delivered, the decision to divert will be influenced by how long ongoing cardiopulmonary resuscitation exists without return of circulation.